




# Maritime Technology and Research

<https://so04.tci-thaijo.org/index.php/MTR>



Research Article

## Species identification of processed sea cucumbers from Malaysian market based on concatenated gene sequences of mitochondrial rRNA genes

Kamarul Rahim Kamarudin<sup>1,\*</sup>, Maryam Mohamed Rehan<sup>2</sup> and Aisyah Mohamed Rehan<sup>3</sup>

<sup>1</sup>Department of Technology and Natural Resources, Faculty of Applied Sciences and Technology, Universiti Tun Hussein Onn Malaysia, Pagoh Campus, Johor Darul Takzim 84600, Malaysia

<sup>2</sup>Department of Food Biotechnology, Faculty of Science and Technology, Universiti Sains Islam Malaysia, Negeri Sembilan Darul Khusus 71800, Malaysia

<sup>3</sup>Department of Biotechnology, Kulliyah of Science, International Islamic University Malaysia, Pahang Darul Makmur 25200, Malaysia

### Article information

Received: September 19, 2018

Revised: December 12, 2018

Accepted: January 10, 2019

### Keywords

12S rRNA gene,  
16S rRNA gene,  
Malaysian markets,  
Processed sea cucumbers,  
Phylogenetic analysis

### Abstract

Species identification of sea cucumbers that have undergone body deformation due to extensive food processing, e.g., beche-de-mer, is difficult, especially with the copresence of cases of unlabelled or mislabelled sea cucumber-based products on the market. Therefore, a study was done to determine the species identities of processed sea cucumbers from selected Malaysian markets using concatenated gene sequences of non-protein-coding 12S and 16S mitochondrial rRNA genes. Phylogenetic analyses, based on the distance-based Neighbor Joining method and character-based methods, i.e., the Maximum Parsimony method, Maximum Likelihood method, and the Bayesian Analysis method, of 47 ingroup sequences, representing 37 processed sea cucumber specimens, 6 reference samples, and 4 additional specimens, suggested the presence of 3 main clusters, i.e., *gamat* family, consisting of genus *Stichopus* and genus *Thelenota*; and *timun laut* family, comprising family Holothuriidae. A number of 3 *gamat* species, i.e., *Stichopus horrens*, *Stichopus vastus*, and *Thelenota anax*, were recorded. Meanwhile, the specimens of *Holothuria (Halodeima) atra*, *Holothuria (Halodeima) edulis*, *Holothuria (Metriatyla) lessoni*, *Holothuria (Mertensiothuria) leucospilota*, and *Holothuria (Metriatyla) scabra* were the 5 *timun laut* species that are grouped under the family Holothuriidae. The outcomes of this study can be utilised by enforcement agencies to monitor and overcome the issues of species substitution and product mislabelling of processed sea cucumber products in Malaysian markets.

*All rights reserved*

## 1. Introduction

The seafood industry is facing a critical issue of species substitution of commercial marine products. There are a number of contributing factors to such issues of intentional product mislabelling; among them are an increase in global seafood consumption, growing international trade, and fluctuations in the food supply and demand of different marine species (Rasmussen &

\*Corresponding author: Department of Technology and Natural Resources, Faculty of Applied Sciences and Technology, Universiti Tun Hussein Onn Malaysia, Pagoh Campus, Johor Darul Takzim 84600, Malaysia  
E-mail address: kamarulr@uthm.edu.my

Morrissey, 2008). According to Rasmussen and Morrissey (2008), economic fraud, health hazards, and illegal trade in protected species could be the serious consequences which result from species substitution. China has commercialised 26 sea cucumber species, placing the country as the second world's top producer (Choo, 2008). However, the mislabelling of 63.6 percent of commercial sea cucumber products from Guangzhou, China, was reported by Wen et al. (2011). The same issue can be observed in Malaysian markets, by which some sea cucumber-based products have been labelled with incorrect species names and are missing manufacturing or packaging details. In fact, Malaysia has been ranked as the world's 4th top producer of commercial sea cucumber, with 19 commercial species (Choo, 2008), and the issue could cause problems in the local economy, health, and conservation sectors.

For years, the microscopic observation of ossicle shapes; the external anatomy of sea cucumber e.g., the presence and shape of tube feet and feeding tentacles; and the internal anatomy, e.g., the types of calcareous rings, have been used for the morphological species identification of sea cucumber. However, even though the morphological characteristics in sea cucumber species identification are important (Dabbagh et al., 2012; Massin et al., 2002), a molecular method, using Deoxyribonucleic Acid (DNA), is required as a confirmation tool, especially for processed sea cucumbers that have undergone shape deformation due to extensive processing. Processed sea cucumbers, including beche-de-mer, are available in the forms of frozen, dried, pickled, and canned products, to mention a few.

Species identification, phylogenetic analyses, and phylogeographical analyses of animals have incorporated mitochondrial DNA (mtDNA) as the most preferred model for molecular genetic studies. Small-subunit mitochondrial ribosomal RNA (rRNA) as part of the mtDNA stores informative resources for phylogenies (Freeman & Herron, 2004). 12S rRNA and 16S rRNA are the 2 components of the small subunit. Neither mitochondrial rRNA gene is a protein-coding gene, since rRNA only produces polypeptides that are used to make up proteins. In fact, the 16S mitochondrial rRNA gene has been frequently used in the molecular genetic analyses of sea cucumber. In terms of species identification of processed sea cucumbers in commercial food products, the PCR-RFLP technique and the Forensically Informative Nucleotide Sequencing (FINS) technique, based on the 16S mitochondrial rRNA gene, have been further developed by Wen et al. (2010) to identify 6 sea cucumber species of the family Stichopodidae. Furthermore, Wen et al. (2011) applied the FINS technique to evaluate the incidence of incorrect labelling of sea cucumbers of the family Holothuriidae. The studies reported the presence of a product mislabeling issue or a species substitution issue in markets. In contrast to 16S mitochondrial rRNA, there has been lack of study on the 12S mitochondrial rRNA gene of sea cucumber to date. Only one study on the 12S mitochondrial rRNA gene of live sea cucumber has been found to date (Clouse et al. 2005). Clouse et al. (2005) summarised that *B. marmorata* and *B. bivittata* should be accepted as 2 separate species, as both species were not sister species, and *B. bivittata* was genetically closer to *B. argus*. In terms of species identification of processed sea cucumbers, Kamarudin et al. (2017) incorporated ossicle shapes and 12S rRNA gene sequences for species identification of *gamat*-based beche-de-mer from Langkawi Island, Kedah, Malaysia. Additionally, Kamarudin et al. (2017) used the 12S mitochondrial rRNA gene to identify the species of sea cucumber specimens from Kudat, Sabah, Malaysia, whereby 3 species were recorded, i.e., *Holothuria scabra*, *Stichopus horrens*, and *Stichopus ocellatus*.


In Malaysia, issues related to species substitution and product mislabelling of sea cucumber-based products can be observed and investigated at some places. Therefore, the aim of this study was to determine the species identity of processed sea cucumber specimens from selected Malaysian markets by using the concatenated gene sequences of non-protein-coding the 12S mitochondrial rRNA gene and the 16S mitochondrial rRNA gene. Since both mitochondrial rRNA genes have been known to be informative, it is believed that their concatenation will give better conclusion on the genetic identity of sea cucumber, due to the interconnection of more informative

sites in a DNA sequence. The genetic identity and relationship of the sea cucumber specimens in this study were determined through the Online Basic Local Alignment Search Tool program for nucleotide (blastn) and phylogenetic analyses based on the distance-based method with clustering algorithm as the tree building strategy, i.e., the Neighbour Joining method, and character-based methods with optimality criterion as the tree building strategy, i.e., the Maximum Parsimony method, Maximum Likelihood method, and the Bayesian Analysis method. A comparison has been made between the findings of the analyses and the manufacturing or packaging details of the sea cucumber specimens. Moreover, this study highlights the issues of intentional species substitution and product labelling of processed sea cucumbers in selected Malaysian markets. The information may be utilised by enforcement agencies to tackle issues pertaining to sea cucumber-based products in Malaysia.

## 2. Materials and methods

### 2.1 Study site and sampling

Kota Kinabalu, Sabah and Kudat, Sabah (East Malaysia, in Borneo Island); Kuantan, Pahang Darul Makmur (East Coast region of Peninsular Malaysia); Langkawi Archipelago, Kedah Darul Aman (North region of Peninsular Malaysia); Nilai, Negeri Sembilan Darul Khusus (South region of Peninsular Malaysia); and Pangkor Archipelago, Perak Darul Ridzuan (West Coast region in the northern part of Peninsular Malaysia) were used as the sampling sites (**Figure 1**). 112 sea cucumber specimens were sampled, including 6 live and fresh specimens of *gamat* species (SHP1-SHP3, *Stichopus horrens*) and *timun laut* species (HLTNP1-HLTNP3, *Holothuria (Mertensiothuria) leucospilota*) from 2 sampling sites in Pangkor Archipelago as the reference samples of fresh sea cucumbers; and 7 dried *gamat*-based beche-de-mer specimens from Kuah, Langkawi Archipelago (LKIG1-LKIG6) as the reference samples of processed sea cucumbers.



**Figure 1** Sampling sites of sea cucumber specimens. Adapted from Kamarudin (2018).

## 2.2 Total genomic DNA extraction

A number of 3 methods of total genomic DNA extraction were used, i.e., the modified cetyl trimethyl ammonium bromide (CTAB) method of Grewe et al. (1993), coupled with the Geneaid Genomic DNA Mini Kit (Blood/Cultured Cell), total genomic DNA extraction using the FavorPrep™ Tissue Genomic DNA Extraction Mini Kit, and total genomic DNA extraction using the DNeasy mericon Food Kit by QIA-GEN. For the third method, homogenized tissue was prepared by using the QIAGEN Tissue-Ruptor for disrupting and homogenising the tissue. One % agarose gel with FloroSafe DNA Stain was used to determine the approximate yield of the total genomic DNA through horizontal gel electrophoresis. The extracts were kept in a -20 °C chest freezer for long-term storage.

## 2.3 Polymerase chain reaction (PCR)

The gene amplification involved 2 methods:

- Twenty five µl PCR reaction volume using the 2x TopTaq Master Mix Kit by QIAGEN;
- Fifty µl PCR reaction volume containing 33.75 µl of sterilized dH<sub>2</sub>O, 5.0 µl of 10X PCR reaction buffer, 3.0 µl of 25 mM magnesium chloride, 2.5 µl of each 5 µM universal primer, 1.0 µl of 10 mM dNTP mix, 2.0 µl of the DNA extract and 0.25 µl of 5 u/µl *Taq* DNA polymerase.

The primer sets for mitochondrial rRNA genes are as follows:

- Primers for 12S mitochondrial rRNA gene (Palumbi et al. (1991), expected length: ~360 bp):  
AB12SA-Lf (forward) 5' – AAA CTG GGA TTA GAT ACC CCA CTA T – 3' (25 bases)  
AB12SB-Hr (reverse) 5' – GAG GGT GAC GGG CGG TGT GT – 3' (20 bases)
- Primers for 16S mitochondrial rRNA gene (Palumbi et al. (1991), expected length: ~650 bp):  
16sar-L (forward) 5' – CGC CTG TTT ATC AAA AAC AT – 3' (20 bases)  
16sbr-H (reverse) 5' – CCG GTC TGA ACT CAG ATC ACG T – 3' (22 bases)

The PCR cycles involved 2 parameter batches:

- 2 min at 95 °C for initial denaturation, 30 s at 95 °C for denaturation, 30 s at optimized temperature for annealing, 45 s at 72 °C for extension, repetition of step 2 - 4 for another 34 - 39 cycles, 5 min at 72 °C for final extension, and forever hold at 4 °C.
- 5 min at 95 °C for initial denaturation, 45 s at 95 °C for denaturation, 90 s at optimized temperature for annealing, 1 min 30 s at 72 °C (60 s/kb; 29 cycles) for extension, 7 min at 72 °C for final extension, and forever hold at 4 °C.

## 2.4 PCR product purification and DNA sequencing

The PCR fragment purification involved 3 types of kit, i.e., QIAquick PCR Purification Kit by QIAGEN (for direct purification of single PCR fragment), Geneaid Gel/PCR DNA Fragments Extraction Kit (for direct purification of single PCR fragment), and QIA-quick Gel Extraction Kit by QIAGEN (for purification of desired PCR fragment from agarose gel). Some of the unpurified PCR products were sent directly to the First BASE Laboratories Sdn Bhd, Seri Kembangan, Selangor Darul Ehsan, Malaysia, as the company also provides PCR product clean-up services.

## 2.5 Phylogenetic analyses

The sequenced PCR products of the mitochondrial DNA genes were displayed using Chromas program version 2.5.1 (Copyright© 1998 - 2016 Technelysium Pty Ltd). The online blastn was used to assign each DNA sequence to a particular sea cucumber species or genus. Prior to the phylogenetic tree reconstruction, ClustalX program version 2.1 (Thompson et al., 1997) was used for multiple sequence alignment of forward reaction sequences. In addition, Molecular Evolutionary Genetics Analysis version 7.0.14 (MEGA7; Kumar et al., 2016) was subsequently used to concatenate

the partial sequences of non-protein-coding 12S and 16S mitochondrial rRNA genes, and also to calculate the number of base substitutions per site from between sequences (i.e., pairwise genetic distance matrix) using the Maximum Composite Likelihood model (Tamura et al., 2004) with the elimination of all positions containing gaps and missing data, and then to reconstruct phylogenetic trees using the Neighbor Joining method (a distance-based method with clustering algorithm as the tree building strategy) and the Maximum Parsimony method (a character-based method with optimality criterion as the tree building strategy). The Modeltest (version 3.7) program (Posada and Crandall, 1998) was used to calculate and find the best model for DNA evolution prior to the reconstruction of Maximum Likelihood phylogenetic trees using the PAUP\* (version 4.0b10) program (Swofford, 1998), with 100 bootstrap replicates. A number of 56 models of DNA substitution were tested in order to choose the model that fitted the data best.

Meanwhile, the reconstructions of consensus Bayesian phylogenetic trees (using the Bayesian Analysis method, a character-based method with optimality criterion as the tree building strategy) were done by using the MrBayes (version 3.1.2) program (Huelsenbeck and Ronquist, 2001). The TreeView (version 1.6.6) program (Page, 1996) and the paint.net 4.0.6 (Final 4.6.5693.28) program (Copyright ©2015 dotPDN LLC, Rick Brewster, and contributors) were used to display and edit the reconstructed phylogenetic trees.

### 3. Results and discussion

**Table 1** indicates the number of base substitutions per site from between concatenated gene sequences of non-protein-coding 12S and 16S mitochondrial rRNA genes. A number of 47 nucleotide sequences and a total of 769 positions were involved in the final dataset. The genetic distance values between specimens that were identified as *S. horrens* ranged from 0 (0 %) to 0.0390 (3.9 %) with average genetic distance of 0.0131 (1.31 %), thus suggesting their status as a single morphospecies, i.e., morphospecies *S. horrens*. Additionally, the genetic distance values between the PKSH1 specimen (from Kudat, Sabah) and other *S. horrens* specimens, including the reference samples of SHP, ranged from 0.0325 (3.25 %) to 0.0390 (3.9 %). Meanwhile, the genetic distance value between *S. vastus* specimens, i.e., PKSO1 (from Kudat, Sabah) and LKIG7, was 0.0185 (1.85 %). The genetic distance values between *T. anax* specimens, i.e., KKS specimens (from Kota Kinabalu, Sabah), ranged from 0.0012 (0.12 %) to 0.0375 (3.75 %), with average genetic distance of 0.0181, thus suggesting their status as a single morphospecies, i.e., morphospecies *T. anax*. Furthermore, the average genetic distance between *Stichopus* specimens was 0.1340 (13.4 %).

**Table 1** Pairwise genetic distance matrix of sequences of concatenated gene sequences of non-protein-coding 12S and 16S mitochondrial rRNA genes of sea cucumber specimens from selected Malaysian markets and other sampling sites, including reference samples and processed specimens.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25
1. HL1																									
2. HLTNP1	0.0278																								
3. HLTNP2	0.0197	0.0105																							
4. HLTNP3	0.0092	0.0211	0.0131																						
5. KKS1	0.3075	0.3135	0.3051	0.3075																					
6. KKS2	0.2967	0.3067	0.2983	0.3007	0.0132																				
7. KKS3	0.2991	0.3091	0.3007	0.3031	0.0145	0.0013																			
8. KKS4	0.2967	0.3067	0.2983	0.3007	0.0145	0.0026	0.0039																		
9. KKS5	0.3468	0.3489	0.3441	0.3468	0.0323	0.0364	0.0379	0.0365																	
10. KKS7	0.3003	0.3063	0.3019	0.3043	0.0185	0.0052	0.0065	0.0078	0.0365																
11. KKS9	0.2967	0.3067	0.2983	0.3007	0.0132	0.0000	0.0013	0.0026	0.0364	0.0052															
12. KPS1	0.1661	0.1661	0.1627	0.1661	0.3091	0.3003	0.3027	0.3003	0.3466	0.3007	0.3003														
13. LKIG1	0.3304	0.3284	0.3300	0.3284	0.2184	0.2086	0.2106	0.2086	0.2469	0.2106	0.2086	0.2596													
14. LKIG2	0.3325	0.3346	0.3362	0.3346	0.2254	0.2121	0.2141	0.2121	0.2543	0.2141	0.2121	0.2672	0.0157												
15. LKIG3	0.3304	0.3325	0.3341	0.3325	0.2201	0.2069	0.2089	0.2069	0.2488	0.2089	0.2069	0.2615	0.0118	0.0039											
16. LKIG4	0.3346	0.3367	0.3383	0.3367	0.2254	0.2138	0.2158	0.2104	0.2525	0.2158	0.2138	0.2653	0.0171	0.0092	0.0052										
17. LKIG5	0.3304	0.3284	0.3300	0.3284	0.2166	0.2069	0.2089	0.2069	0.2451	0.2089	0.2069	0.2596	0.0039	0.0118	0.0078	0.0131									
18. LKIG6	0.3284	0.3304	0.3321	0.3304	0.2184	0.2051	0.2071	0.2051	0.2469	0.2071	0.2051	0.2615	0.0052	0.0105	0.0065	0.0118	0.0013								
19. LKIG7	0.3518	0.3518	0.3492	0.3455	0.2093	0.1997	0.2017	0.1997	0.2356	0.2017	0.1997	0.2811	0.0497	0.0470	0.0429	0.0429	0.0456	0.0470							
20. PFK1	0.1480	0.1576	0.1542	0.1544	0.3152	0.3043	0.3067	0.3043	0.3551	0.3039	0.3043	0.1273	0.3075	0.3115	0.3055	0.3095	0.3075	0.3055	0.3279						
21. PFK11	0.1463	0.1558	0.1525	0.1526	0.3177	0.3067	0.3091	0.3067	0.3578	0.3063	0.3067	0.1256	0.3099	0.3140	0.3079	0.3119	0.3099	0.3079	0.3254	0.0013					
22. PFK12	0.1479	0.1542	0.1541	0.1542	0.3197	0.3087	0.3112	0.3087	0.3556	0.3043	0.3087	0.1271	0.3119	0.3160	0.3099	0.3140	0.3119	0.3099	0.3274	0.0026	0.0013				
23. PFK13	0.1774	0.1823	0.1821	0.1823	0.3442	0.3442	0.3468	0.3442	0.3864	0.3395	0.3442	0.1543	0.3432	0.3496	0.3432	0.3475	0.3432	0.3432	0.3596	0.0294	0.0281	0.0267			
24. PFK14	0.1514	0.1609	0.1576	0.1577	0.3218	0.3107	0.3132	0.3107	0.3621	0.3103	0.3107	0.1305	0.3140	0.3180	0.3119	0.3160	0.3140	0.3119	0.3325	0.0039	0.0039	0.0052	0.0322		
25. PFK15	0.1479	0.1542	0.1541	0.1542	0.3197	0.3087	0.3112	0.3087	0.3556	0.3043	0.3087	0.1271	0.3119	0.3160	0.3099	0.3140	0.3119	0.3099	0.3274	0.0026	0.0013	0.0000	0.0267	0.0052	



	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25
26. PFKK16	0.1514	0.1609	0.1576	0.1577	0.1377	0.3067	0.3091	0.3067	0.3578	0.3063	0.3067	0.1305	0.3099	0.3140	0.3079	0.3119	0.3099	0.3079	0.3304	0.0026	0.0039	0.0052	0.0322	0.0065	0.0052
27. PFKK2	0.1479	0.1574	0.1541	0.1542	0.3156	0.3047	0.3071	0.3047	0.3556	0.3043	0.3047	0.1256	0.3099	0.3140	0.3079	0.3119	0.3099	0.3079	0.3254	0.0026	0.0013	0.0026	0.0294	0.0052	0.0026
28. PFKK3	0.1465	0.1560	0.1526	0.1528	0.3132	0.3023	0.3047	0.3023	0.3530	0.3019	0.3023	0.1273	0.3095	0.3136	0.3075	0.3115	0.3095	0.3075	0.3300	0.0013	0.0026	0.0039	0.0308	0.0052	0.0039
29. PFKK4	0.1465	0.1560	0.1526	0.1528	0.3132	0.3023	0.3047	0.3023	0.3530	0.3019	0.3023	0.1273	0.3095	0.3136	0.3075	0.3115	0.3095	0.3075	0.3300	0.0013	0.0026	0.0039	0.0308	0.0052	0.0039
30. PFKK5	0.1465	0.1560	0.1526	0.1528	0.3132	0.3023	0.3047	0.3023	0.3530	0.3019	0.3023	0.1273	0.3095	0.3136	0.3075	0.3115	0.3095	0.3075	0.3300	0.0013	0.0026	0.0039	0.0308	0.0052	0.0039
31. PFKK6	0.1683	0.1715	0.1681	0.1666	0.3313	0.3172	0.3197	0.3193	0.3692	0.3176	0.3172	0.1486	0.2959	0.2939	0.2919	0.2939	0.2959	0.2939	0.3070	0.1100	0.1084	0.1099	0.1372	0.1131	0.1099
32. PFKK7	0.1463	0.1558	0.1525	0.1526	0.3177	0.3067	0.3091	0.3067	0.3578	0.3063	0.3067	0.1256	0.3099	0.3140	0.3079	0.3119	0.3099	0.3079	0.3254	0.0013	0.0000	0.0013	0.0281	0.0039	0.0013
33. PFKK8	0.1905	0.1989	0.1970	0.1972	0.3638	0.3520	0.3547	0.3520	0.4076	0.3515	0.3520	0.1646	0.3491	0.3512	0.3448	0.3491	0.3491	0.3470	0.3709	0.0378	0.0393	0.0406	0.0696	0.0420	0.0406
34. PFKK9	0.1809	0.1859	0.1856	0.1858	0.3606	0.3563	0.3589	0.3563	0.3949	0.3515	0.3563	0.1646	0.3489	0.3532	0.3489	0.3531	0.3489	0.3489	0.3654	0.0364	0.0350	0.0336	0.0199	0.0392	0.0336
35. PK51	0.1729	0.1761	0.1727	0.1761	0.3156	0.2999	0.3023	0.3019	0.3525	0.2983	0.2999	0.0158	0.2556	0.2594	0.2537	0.2593	0.2556	0.2537	0.2730	0.1305	0.1288	0.1304	0.1594	0.1338	0.1304
36. PK52	0.1712	0.1778	0.1743	0.1745	0.3156	0.2999	0.3023	0.3019	0.3525	0.2983	0.2999	0.0171	0.2574	0.2612	0.2556	0.2612	0.2574	0.2556	0.2711	0.1321	0.1304	0.1320	0.1610	0.1354	0.1320
37. PK521	0.1834	0.1867	0.1832	0.1867	0.3284	0.3124	0.3148	0.3144	0.3660	0.3108	0.3124	0.0265	0.2673	0.2711	0.2654	0.2711	0.2673	0.2654	0.2850	0.1436	0.1418	0.1434	0.1731	0.1469	0.1434
38. PK53	0.1729	0.1761	0.1727	0.1761	0.3156	0.2999	0.3023	0.3019	0.3503	0.2983	0.2999	0.0184	0.2574	0.2612	0.2556	0.2612	0.2574	0.2556	0.2749	0.1274	0.1257	0.1273	0.1561	0.1307	0.1273
39. PK54	0.1745	0.1778	0.1743	0.1778	0.3177	0.3019	0.3043	0.3039	0.3525	0.3003	0.3019	0.0197	0.2593	0.2632	0.2575	0.2631	0.2593	0.2575	0.2769	0.1258	0.1242	0.1257	0.1545	0.1291	0.1257
40. PK5H1	0.3263	0.3242	0.3258	0.3242	0.2219	0.2086	0.2106	0.2086	0.2506	0.2106	0.2086	0.2673	0.0331	0.0304	0.0264	0.0318	0.0291	0.0278	0.0484	0.3075	0.3099	0.3120	0.3454	0.3140	0.3120
41. PK5O1	0.3383	0.3404	0.3378	0.3362	0.2111	0.1980	0.1999	0.1980	0.2374	0.1999	0.1980	0.2733	0.0414	0.0387	0.0346	0.0400	0.0373	0.0360	0.0158	0.3148	0.3123	0.3144	0.3479	0.3193	0.3144
42. PM1	0.1479	0.1574	0.1541	0.1542	0.3156	0.3047	0.3071	0.3047	0.3556	0.3043	0.3047	0.1256	0.3099	0.3140	0.3079	0.3119	0.3099	0.3079	0.3254	0.0026	0.0013	0.0026	0.0294	0.0052	0.0026
43. PM3	0.1683	0.1715	0.1681	0.1666	0.3313	0.3172	0.3197	0.3193	0.3692	0.3176	0.3172	0.1486	0.2959	0.2939	0.2919	0.2939	0.2959	0.2939	0.3070	0.1100	0.1084	0.1099	0.1372	0.1131	0.1099
44. PM4	0.2921	0.2970	0.2859	0.2901	0.4813	0.4828	0.4862	0.4828	0.5069	0.4820	0.4828	0.2754	0.4357	0.4309	0.4309	0.4309	0.4357	0.4333	0.4523	0.2244	0.2224	0.2242	0.2420	0.2283	0.2242
45. SHP1	0.3304	0.3283	0.3300	0.3283	0.2201	0.2104	0.2123	0.2103	0.2488	0.2123	0.2104	0.2615	0.0118	0.0039	0.0026	0.0078	0.0078	0.0092	0.0429	0.3095	0.3119	0.3140	0.3453	0.3160	0.3140
46. SHP2	0.3304	0.3283	0.3300	0.3283	0.2201	0.2104	0.2123	0.2103	0.2488	0.2123	0.2104	0.2615	0.0118	0.0039	0.0026	0.0078	0.0078	0.0092	0.0429	0.3095	0.3119	0.3140	0.3453	0.3160	0.3140
47. SHP3	0.3283	0.3304	0.3320	0.3304	0.2219	0.2086	0.2106	0.2086	0.2506	0.2106	0.2086	0.2634	0.0131	0.0026	0.0013	0.0065	0.0092	0.0078	0.0442	0.3075	0.3099	0.3119	0.3453	0.3140	0.3119

	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47
26. PFKK16																						
27. PFKK2	0.0052																					
28. PFKK3	0.0039	0.0039																				
29. PFKK4	0.0039	0.0039	0.0000																			
30. PFKK5	0.0039	0.0039	0.0000	0.0000																		
31. PFKK6	0.1131	0.1084	0.1085	0.1085	0.1085																	
32. PFKK7	0.0039	0.0013	0.0026	0.0026	0.0026	0.1084																
33. PFKK8	0.0406	0.0393	0.0365	0.0365	0.0365	0.1477	0.0393															
34. PFKK9	0.0392	0.0363	0.0378	0.0378	0.0378	0.1458	0.0350	0.0771														
35. PKS1	0.1338	0.1288	0.1305	0.1305	0.1305	0.1422	0.1288	0.1681	0.1698													
36. PKS2	0.1354	0.1304	0.1321	0.1321	0.1321	0.1438	0.1304	0.1698	0.1714	0.0013												
37. PKS21	0.1469	0.1418	0.1436	0.1436	0.1436	0.1522	0.1418	0.1804	0.1816	0.0105	0.0118											
38. PKS3	0.1307	0.1257	0.1274	0.1274	0.1274	0.1390	0.1257	0.1648	0.1664	0.0026	0.0039	0.0131										
39. PKS4	0.1291	0.1242	0.1258	0.1258	0.1258	0.1375	0.1242	0.1632	0.1648	0.0039	0.0052	0.0145	0.0013									
40. PKS41	0.3099	0.3099	0.3095	0.3095	0.3095	0.2880	0.3099	0.3491	0.3510	0.2556	0.2575	0.2673	0.2575	0.2594								
41. PKS01	0.3172	0.3123	0.3168	0.3168	0.3168	0.2962	0.3123	0.3548	0.3536	0.2616	0.2634	0.2734	0.2635	0.2654	0.0373							
42. PM1	0.0052	0.0000	0.0039	0.0039	0.0039	0.1084	0.0013	0.0393	0.0363	0.1288	0.1304	0.1418	0.1257	0.1242	0.3099	0.3123						
43. PM3	0.1131	0.1084	0.1085	0.1085	0.1085	0.0000	0.1084	0.1477	0.1458	0.1422	0.1438	0.1522	0.1390	0.1375	0.2880	0.2962	0.1084					
44. PM4	0.2283	0.2224	0.2226	0.2226	0.2226	0.1245	0.2224	0.2716	0.2527	0.2758	0.2777	0.2865	0.2739	0.2720	0.4285	0.4388	0.2224	0.1245				
45. SHP1	0.3119	0.3119	0.3115	0.3115	0.3115	0.2959	0.3119	0.3491	0.3510	0.2575	0.2593	0.2692	0.2594	0.2612	0.0291	0.0373	0.3119	0.2959	0.4357			
46. SHP2	0.3119	0.3119	0.3115	0.3115	0.3115	0.2959	0.3119	0.3491	0.3510	0.2575	0.2593	0.2692	0.2594	0.2612	0.0291	0.0373	0.3119	0.2959	0.4357	0.0000		
47. SHP3	0.3099	0.3099	0.3095	0.3095	0.3095	0.2939	0.3099	0.3470	0.3510	0.2556	0.2575	0.2673	0.2575	0.2594	0.0278	0.0360	0.3099	0.2939	0.4333	0.0013	0.0013	




With regard to the *timun laut* specimens, the average genetic distance value between *Holothuria* specimens, including the reference samples of HLTNP, was 0.1112 (11.12 %). Moreover, the genetic distance values between *H. leucospilota* specimens (HLTNP specimens and the HL1 specimen from Pangkor Archipelago, Perak) ranged from 0.0086 (0.86 %) to 0.0277 (2.77 %), with an average genetic distance of 0.0169 (1.69 %). The genetic distance values between the HL1 specimen (from Pangkor Archipelago, Perak) and other *H. leucospilota* specimens ranged from 0.0086 (0.86 %) to 0.0277 (2.77 %). Furthermore, the genetic distance values between *H. scabra* specimens (PKS specimens from Kudat, Sabah) ranged from 0.0013 (0.13 %) to 0.0140 (1.4 %), with an average genetic distance of 0.0066 (0.66 %), while the genetic distance values between the *H. lessoni* specimen, i.e., KPTS1 (from Kuantan, Pahang) and *H. scabra* specimens ranged from 0.0179 (1.79 %) to 0.0284 (2.84 %). As for the *H. atra* specimens (PFFK6 specimen from Kota Kinabalu, Sabah, and specimens of PM3 and PM4 from Manukan Island, Sabah), the average genetic distance was 0.0835 (8.35 %), ranging from 0 (0 %) to 0.1253 (12.53 %), with the genetic distance value between the PFFK6 specimen and the PM3 specimen being 0 (0 %). The average genetic distance between *H. edulis* specimens (PFFK specimens from Kota Kinabalu, Sabah, and PM1 specimen from Manukan Island, Sabah), excluding the PFFK6 specimen, was 0.0293 (2.93 %), ranging from 0 (0 %) to 0.1539 (15.39 %), thus suggesting their status as a single morphospecies, i.e., morphospecies *H. edulis*. The genetic distance values between the PM1 specimen and PFFK specimens, excluding the PFFK6 specimen, ranged from 0 (0 %) to 0.1182 (11.82 %).


With regard to the Neighbor Joining analyses using concatenated gene sequences of the non-protein-coding 12S and 16S mitochondrial rRNA genes, the optimal tree with the sum of branch length = 7.50559484 is shown in **Figure 2**. The percentage of replicate trees in which the associated taxa clustered together in the boot-strap test (i.e., 1000 replicates) are shown next to the branches (Felsenstein, 1985). The tree was drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic tree. The evolutionary distances were computed using the Maximum Composite Likelihood method (Tamura et al., 2004), and are in the units of the number of base substitutions per site. All positions containing gaps and missing data were eliminated. A number of 48 taxa, consisting of 47 ingroup taxa and one outgroup taxon, and 749 characters, representing aligned base positions (after multiple alignment), were involved in the phylogenetic analyses of the sequences of the concatenated gene sequences of the non-protein-coding 12S and 16S mitochondrial rRNA genes using the Neighbor Joining method. The outgroup taxon was an individual of *Peniagone* sp., a deep-sea swimming sea cucumber species, with GenBank Accession No. KF915304.

**Figure 2** illustrates the presence of 2 main groups of the specimens: the *timun laut* family, with 47 % bootstrap support, and a few clusters representing family Stichopodidae (the *gamat* family). A number of 3 *gamat* species, i.e., *S. horrens*, *S. vastus*, and *T. anax*, represented the family Stichopodidae. *T. anax* was divided into 2 clusters, and *S. horrens* cluster was supported by 96 % bootstrap value. Nonetheless, the specimens of *S. vastus* were also grouped into *S. horrens*, due to their close genetic relationship. As mentioned earlier, the genetic distance value between *S. vastus* specimens, i.e. PKSO1 (from Kudat, Sabah) and LKIG7, was low, i.e., 0.0185 (1.85 %). *S. vastus* cluster was supported with 97 % bootstrap value. In addition, the specimens of PFFK, excluding the PFFK6 specimen, were clustered as *H. edulis*, with 91 % bootstrap support. Purcell et al. (2012) recorded a processed *H. lessoni* that is similar to the dried *tip-sum* (KPTS1); therefore, the specimen was regarded as *H. lessoni*. In terms of the species status of PM specimens, the analyses identified the PM1 specimen as *H. edulis*, and the specimens of PM3 and PM4 as *H. atra*. Therefore, the specimens of *H. leucospilota* that formed a cluster with 92 % bootstrap value, the specimens of *H. edulis* that formed a cluster with 91 % bootstrap value, the specimens of *H. scabra* that formed a cluster with 66 % bootstrap value, the dried specimen of *H. lessoni* (KPTS1), and the specimens of *H. atra* that formed a cluster with 79 % bootstrap value were the 5 *timun laut* species that are grouped under the family Holothuriidae, with 47 % bootstrap support. *H. atra* was genetically

closer to *H. edulis*, with 68 % bootstrap support. Furthermore, the subgenus *Mertensiothuria*, represented by the specimens of *H. leucospilota*, was genetically closer to the subgenus *Halodeima*, represented by the specimens of *H. atra* and *H. edulis*, with 46 % bootstrap value, thus supporting their taxonomic classification.



**Figure 2** Topology of 50 % majority-rule consensus tree of Neighbor Joining of sea cucumber specimens from selected Malaysian markets and other sampling sites, including the reference samples and processed specimens inferred from concatenated gene sequences of non-protein-coding 12S and 16S mitochondrial rRNA genes using MEGA7 program (Kumar et al., 2016), with 1000 bootstrap replicates. Numbers at nodes indicate the bootstrap values in percentage (%).



**Figure 3** Topology of 50 % majority-rule consensus tree of Maximum Parsimony of sea cucumber specimens from selected Malaysian markets and other sampling sites, including the reference samples and processed specimens inferred from concatenated gene sequences of non-protein-coding 12S and 16S mitochondrial rRNA genes using MEGA7 program (Kumar et al., 2016), with 1000 bootstrap replicates. Numbers at nodes indicate the bootstrap values in percentage (%).

As for the Maximum Parsimony analyses using concatenated gene sequences of the non-protein-coding 12S and 16S mitochondrial rRNA genes, the bootstrap consensus tree inferred from 1000 replicates was taken to represent the evolutionary history of the taxa analyzed (Felsenstein, 1985). Branches corresponding to partitions reproduced in less than 50 % bootstrap replicates were collapsed. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) are shown next to the branches (Felsenstein, 1985). The Maximum Parsimony tree was obtained using the Subtree-Pruning-Regrafting (SPR) algorithm (Nei & Kumar, 2000) with search level 1, in which the initial trees were obtained by the random addition of sequences (10 replicates). All positions containing gaps and missing data were eliminated. Additionally, a number of 48 taxa, consisting of 47 ingroup taxa and one outgroup taxon, and 749 characters, representing aligned base positions (after multiple alignment), were involved in the phylogenetic analyses.

**Figure 3** illustrates the presence of 2 main clusters of the specimens: family Stichopodidae (the *gamat* family) and the *timun laut* family, with 90 % bootstrap support and 99 % bootstrap support, respectively. A number of 3 *gamat* species, i.e., *S. horrens*, *S. vastus*, and *T. anax*, were clustered under the family Stichopodidae. *T. anax* cluster was supported by 99 % bootstrap value, *S. horrens* cluster was supported by 76 % bootstrap value, and *S. vastus* cluster was supported by 93 % bootstrap value. *S. vastus* was genetically closer to *S. horrens*, with 99 % bootstrap support. Furthermore, the specimens of *H. leucospilota* that formed a cluster with 99 % bootstrap value, the specimens of *H. edulis* that formed a cluster with 99 % bootstrap value, the specimens of *H. scabra* that formed a cluster with 88 % bootstrap value, the dried specimen of *H. lessoni* (KPTS1), and the specimens of *H. atra* that formed a cluster with 99 % bootstrap value were the 5 *timun laut* species that is grouped under the family Holothuriidae, with 98 % bootstrap support. *H. atra* was genetically closer to *H. edulis*, with 51 % bootstrap support. Likewise, in the Neighbor Joining analyses, the subgenus *Mertensiothuria*, represented by the specimens of *H. leucospilota*, was genetically closer to the subgenus *Halodeima*, represented by the specimens of *H. atra* and *H. edulis*, with 93 % bootstrap value.

Pertaining to the Maximum Likelihood analyses using the concatenated gene sequences of non-protein-coding 12S and 16S mitochondrial rRNA genes, the Modeltest (version 3.7) program suggested the general time-reversible (GTR) model (Tavaré, 1986), with the rate variation among sites (+G) (i.e., GTR+G) as the best model of DNA substitution, based on the Akaike Information Criterion (AIC); and the Tamura and Nei (TrN) model (Tamura & Nei, 1993), with the rate variation among sites (+G) (i.e., TrN+G) as the best model of DNA substitution, based on the Hierarchical Likelihood Ratio Tests (hLRTs). The GTR model and the TrN model were based on unequal base frequencies. According to Posada and Buckley (2004), the AIC and Bayesian Analyses are good for the evaluation of model selection uncertainty, capable of comparing multiple nested or non-nested models at once, and allow for the use of all available models for the estimation of phylogenies and model parameters. Hence, between the AIC and the hLRTs used in the Modeltest program, the model of DNA substitution suggested by the AIC is better. The GTR+G model suggested by the AIC was chosen for the Maximum Likelihood tree reconstruction (Lset Base=(0.3343 0.2070 0.1869) Nst=6 Rmat=(2.4444 4.2221 1.8990 1.0703 7.6112) Rates=gamma Shape=0.7426 Pinvar=0)).


A number of 48 taxa, consisting of 47 ingroup taxa and one outgroup taxon, and 859 characters, representing aligned base positions (after multiple alignment), were involved in the phylogenetic analyses. Overall, the base frequencies were unequal (i.e. Adenine (A) = 33.43 %, Cytosine (C) = 20.70 %, Guanine (G) = 18.69 %, and Thymine (T) = 27.18 %), thus supporting the selection of GTR+G model by the AIC as the best model. In addition, **Figure 4** illustrates the presence of 2 clusters of family Stichopodidae (the *gamat* family), i.e., genus *Stichopus* cluster, with 86 % bootstrap support, and genus *Thelenota* cluster, with 66 % bootstrap support; and the *timun laut* family, with 59 % bootstrap support. A number of 3 *gamat* species, i.e., *S. horrens*, *S.*

*vastus*, and *T. anax* under the family Stichopodidae, were identified. *T. anax* cluster was supported by 66 % bootstrap value, *S. horrens* cluster was supported by 62 % boot-strap value, and *S. vastus* cluster was supported by 94 % bootstrap value. *S. vastus* was genetically closer to *S. horrens*, with 86 % bootstrap support. Additionally, the specimens of *H. leucospilota* that formed a cluster with 91 % bootstrap value, the specimens of *H. edulis* that formed a cluster with 90 % bootstrap value, the specimens of *H. scabra* (with the inclusion of the dried specimen of *H. lessoni* (KPTS1)) with 90 % bootstrap value, and the specimens of *H. atra* that formed a cluster with 98 % bootstrap value were the 5 *timun laut* species that is grouped under the family Holothuriidae, with 59 % bootstrap support.

Furthermore, the phylogenetic analysis of the Bayesian Analysis method was ended when the standard deviation of split frequencies was below 0.01. Accordingly, for the Bayesian Analysis analyses of the concatenated gene sequences of non-protein-coding 12S and 16S mitochondrial rRNA genes, the standard deviation of split frequencies was 0.007697 at 1,780,000 generations. **Figure 5** illustrates the presence of 3 main groups of the specimens: family Stichopodidae (the *gamat* family), with 66 % posterior probability, and 2 subclusters of the *timun laut* family. A number of 3 *gamat* species, i.e., *S. horrens*, *S. vastus*, and *T. anax*, were clustered under the family Stichopodidae. *T. anax* cluster was supported by 87 % posterior probability, *S. horrens* cluster was supported by 50 % posterior probability, and *S. vastus* cluster was supported by 99 % posterior probability.


The specimens of *H. leucospilota* that formed a cluster with 98 % posterior probability, the specimens of *H. edulis* that formed a cluster with 95 % posterior probability, and the specimens of *H. atra* that formed a cluster with 89 % posterior probability were the 3 *timun laut* species that formed one of the subclusters of the *timun laut* family, with 65 % posterior probability. The specimens of *H. scabra* that formed a cluster with 94 % posterior probability and the dried specimen of *H. lessoni* (KPTS1) were the 2 *timun laut* species that formed the other subcluster of the *timun laut* family, with 90 % posterior probability. Moreover, *H. edulis* was closer to *H. atra*, with 75 % posterior probability, and *H. scabra* was closer to *H. lessoni*, with 90 % posterior probability. Likewise, in the Neighbor Joining analyses and the Maximum Parsimony analyses, the subgenus *Mertensiothuria*, represented by the specimens of *H. leucospilota*, was genetically closer to the subgenus *Halodeima*, represented by the specimens of *H. atra* and *H. edulis*, with 65 % posterior probability.

In addition, the Maximum Likelihood tree (**Figure 4**) and the Bayesian Analysis tree (**Figure 5**) supported that *H. scabra* was genetically closer to *H. lessoni*, with 90 % bootstrap value/posterior probability, thus supporting their taxonomic classification as from the subgenus *Metriatyla*. Except for the Bayesian Analysis tree, the other phylogenetic trees showed the clustering of *timun laut* specimens into clusters, with 47 - 99 % bootstrap values, thus suggesting the formation of the *timun laut* group. In summary, 8 sea cucumber species were recorded in this study, including 5 *timun laut* species, i.e., *H. leucospilota*, *H. atra*, *H. edulis*, *H. scabra*, and *H. lessoni*; and 3 *gamat* species, i.e., *S. horrens*, *S. vastus*, and *T. anax*. *H. leucospilota*, *H. atra*, *H. edulis*, *H. scabra*, *S. horrens*, and *T. anax* were commercial Malaysian sea cucumber species (Choo, 2008). Nevertheless, *H. lessoni* and *S. vastus* were not listed as commercial Malaysian sea cucumber species. Among the species recorded in this study, 2 *timun laut* species were included in the International Union for Conservation of Nature (IUCN) Red List for aspidochirotid holothuroids, whereby *H. lessoni* and *H. scabra* were regarded as “endangered, or at a high risk of extinction” (Conand et al., 2014). Apart from that, the outcomes of this study provide better information on the level of species substitution and product mislabeling issues of processed sea cucumbers in Malaysian markets, which may subsequently assist enforcement agencies to monitor and overcome these issues through the introduction of mtDNA sequencing techniques.



**Figure 4** Topology of 50 % majority-rule consensus tree of Maximum Likelihood of sea cucumber specimens from selected Malaysian markets and other sampling sites, including the reference samples and processed specimens inferred from concatenated gene sequences of non-protein-coding 12S and 16S mitochondrial rRNA genes using PAUP\* (version 4.0b10) program (Swofford, 1998), with 100 bootstrap replicates. Numbers at nodes indicate the bootstrap values in percentage (%).





**Figure 5** Topology of consensus Bayesian Analysis tree of sea cucumber specimens from selected Malaysian markets and other sampling sites, including the reference samples and processed specimens inferred from concatenated gene sequences of non-protein-coding 12S and 16S mitochondrial rRNA genes using MrBayes (version 3.1.2) program (Huelsenbeck & Ronquist, 2001), with the addition of all compatible groups to the tree. Numbers at nodes indicate the posterior probabilities of clades in percentage (%).

#### 4. Conclusions

In conclusion, the phylogenetic trees based on the distance-based method with clustering algorithm as the tree building strategy, i.e., the Neighbor Joining method, and character-based methods with optimality criterion as the tree building strategy, i.e., the Maximum Parsimony method, the Maximum Likelihood method, and the Bayesian Analysis method, suggested the presence of 3 main clusters of the specimens, i.e., the *gamat* family, consisting of genus *Stichopus* and genus *Thelenota*; and the *timun laut* family, comprising family Holothuriidae. Three *gamat* species, i.e., *S. horrens*, *S. vastus*, and *T. anax*; and 5 *timun laut* species, i.e., *H. atra*, *H. edulis*, *H. lessoni*, *H. leucospilota*, and *H. scabra* were recorded. This study also highlights the presence of issues of intentional species substitution or product mislabeling due to the observation of unlabelled products in the selected Malaysian markets. The outcomes of this study may assist enforcement agencies to monitor and address these said issues.

## Acknowledgements

We would like to thank Mrs. Siti Nabilah Bt Mohd Rusly, Mr. Muhammad Izzat B. Redzuan, and Ms. Sarina Irma Binti Saidon; all members of Food Biotechnology program, Faculty of Science and Technology (FST), Universiti Sains Islam Malaysia (USIM), Nilai Campus; all reviewers of this paper; and the Malaysian Ministry of Higher Education (MOHE) for input and financial assistance. Research Acculturation Grant Scheme (RAGS) Phase 1/2014 from the Ministry of Education (MOE) - Ref: USIM/RAGS/FST/36/50414 and the Fundamental Research Grant Scheme (FRGS) Phase 1/2015 from the Department of Higher Education, MOE - Ref: USIM/FRGS/FST/32/51515 were used to fund this research. Visit Malaysian Sea Cucumber Database at <http://sites.google.com/site/malaysianseacucumber/> for more details.

## References

- Choo, P. S. (2008). Population status, fisheries and trade of sea cucumbers in Asia. In V. Toral-Granda, A. Lovatelli & M. Vasconcellos (Eds.). *Sea cucumbers. A global review of fisheries and trade* (pp. 81-118). FAO Fisheries and Aquaculture Technical Paper. No. 516., Rome.
- Clouse, R., Janies, D., & Kerr, A. M. (2005). Resurrection of *Bohadschia bivittata* from *B. marmorata* (Holothuroidea: Holothuriidae) based on behavioral, morphological, and mitochondrial DNA evidence. *Zoology*, 108(1), 27-39. doi:10.1016/j.zool.2004.07.007
- Conand, C., Polidoro, B., Mercier, A., Gamboa, R., Hamel, J. F., & Purcell, S. (2014). The IUCN Red List assessment of aspidochirotid sea cucumbers and its implications. *SPC Beche-de-mer Information Bulletin*, 34(5), 3-7.
- Dabbagh, A. R., Keshavarz, M., Mohammadikia, D., Afkhami, M., & Nateghi, S. A. (2012). *Holothuria scabra* (Holothuroidea: Aspidochirotida): First record of a highly valued sea cucumber, in the Persian Gulf, Iran. *Marine Biodiversity Records*, 5(8), e69. doi:10.1017/S1755267212000620
- Freeman, S., & Herron, J. C. (2004). *Evolutionary Analysis*. 3<sup>rd</sup> ed. Upper Saddle River, NJ: Pearson/Prentice Hall.
- Huelsenbeck, J. P., & Ronquist, F. (2001). MRBAYES: Bayesian inference of phylogeny. *Bioinformatics*, 17(8), 754-755. doi:10.1093/bioinformatics/17.8.754
- Kamarudin, K. R. (2018). *Species Identification, Molecular Phylogenetic and Genetic Diversity of Sea Cucumber (Phylum Echinodermata: Class Holothuroidea) in Malaysia*. Ph.D. Thesis, Universiti Sains Islam Malaysia, Nilai, Negeri Sembilan, Malaysia.
- Kamarudin, K. R., Mohamed, M. R., & Bahaman, N. A. (2017). Morphological and molecular identification of ossicle shapes in sea cucumber species *Holothuria scabra*, *Stichopus horrens* and *Stichopus ocellatus* from Kudat, Sabah, Malaysia. *Pertanika Journal of Tropical Agricultural Science*, 40(1), 161-172.
- Kamarudin, K. R., Mohamed, M. R., Mohd, H. N., Ramly, N. Z., & Mohamed, A. R. (2017). Comparison of ossicle shape and 12S rRNA gene sequencing techniques for species identification of *gamat*-based beche-de-mer from Langkawi Archipelago, Kedah Darul Aman. *Pertanika Journal of Tropical Agricultural Science*, 40(2), 91-104.
- Kumar, S., Stecher, G., & Tamura, K. (2016). MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution*, 33(7), 1870-1874. doi:10.1093/molbev/msw054
- Massin, C., Zulfigar, Y., Hwai, T. S., & Boss, S. Z. (2002). The genus *Stichopus* (Echinodermata: Holothuroidea) from the Johore Marine Park (Malaysia) with the description of two new species. *Bulletin de l'Institut royal des Sciences naturelles de Belgique*, 72, 73-99.
- Page, R. D. M. (1996). TREEVIEW: An application to display phylogenetic trees on personal computers. *Bioinformatics*, 12(4), 357-358. doi:10.1093/bioinformatics/12.4.357

- Palumbi, S., Martin, A., Romano, S., McMillan, W. O., Stice, L., & Grabowski, G. (1991). *The Simple Fool's Guide to PCR*. Department of Zoology and Kewalo Marine Laboratory, Universiti of Hawaii, Honolulu.
- Posada, D., & Buckley, T. R. (2004). Model selection and model averaging in phylogenetics: Advantages of akaike information criterion and bayesian approaches over likelihood ratio tests. *Systematic Biology*, 53(5), 793-808. doi:10.1080/10635150490522304
- Posada, D., & Crandall, K. A. (1998). Modeltest: Testing the model of DNA substitution. *Bioinformatics*, 14(9), 817-818. doi:10.1093/bioinformatics/14.9.817
- Rasmussen, R. S., & Morrissey, M. T. (2008). DNA-based methods for the identification of commercial fish and seafood species. *Comprehensive Reviews in Food Science and Food Safety*, 7(3), 280-295. doi:10.1111/j.1541-4337.2008.00046.x
- Swofford, D. L. (1998). "PAUP", Phylogenetic Analysis Using Parsimony (\*and other methods), version 4. Sinauer Associates, Sunderland, MA.
- Tamura, K., Nei, M., & Kumar, S. (2004). Prospects for inferring very large phylogenies by using the neighbor-joining method. *Proceedings of the National Academy of Sciences of the United States of America*, 101(30), 11030-11035. doi:10.1073/pnas.0404206101
- Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F., & Higgins, D. G. (1997). The ClustalX windows interface: Flexible strategies for multiple sequence alignment aided by the quality analysis tools. *Nucleic Acids Research*, 25(24), 4876-4882. doi:10.1093/nar/25.24.4876
- Wen, J., Hu, C., Zhang, L., & Fan, S. (2011). Genetic identification of global commercial sea cucumber species on the basis of mitochondrial DNA sequences. *Food Control*, 22(1), 72-77. doi:10.1016/j.foodcont.2010.06.010
- Wen, J., Hu, C., Zhang, L., Luo, P., Zhao, Z., Fan, S., & Su, T. (2010). The application of PCR-RFLP and FINS for species identification used in sea cucumbers (Aspidochirotida: Stichopodidae) products from the market. *Food Control*, 21(4), 403-407. doi:10.1016/j.foodcont.2009.06.014